

ON EXTENSIONS OF GROUP WITH INFINITE CONJUGACY CLASSES, I

JEAN-PHILIPPE PRÉAUX^{1 2}

ABSTRACT. We give a characterization of the group property of being with infinite conjugacy classes (or *icc*, *i.e.* $\neq 1$ and of which all conjugacy classes beside 1 are infinite) for extensions of abelian, centerless, *icc*, or word hyperbolic groups.

INTRODUCTION

A group is said to be with *infinite conjugacy classes* (or *icc*) if it is non trivial, and if all its conjugacy classes beside $\{1\}$ are infinite. This property is motivated by the theory of Von Neumann algebra, since for any group Γ , a necessary and sufficient condition for its Von Neumann algebra $W_\lambda^*(\Gamma)$ to be a type $II - 1$ factor is that Γ be *icc* (cf. [ROIV]).

The property of being *icc* has been characterized in several classes of groups : 3-manifolds and $PD(3)$ groups in [HP], groups acting on Bass-Serre trees in [Co], wreath products, semi-direct products and finite extensions in [P1, P2, P3]. We will focus here on groups defined by some specific extensions of a group, namely extensions of abelian groups, of centerless groups, of *icc* groups, or of word hyperbolic groups. We give a necessary and sufficient condition for those groups to satisfy the *icc* property.

We first consider the case of extensions of abelian groups. In a group G , $FC(G)$ denotes the unions of finite conjugacy classes in G ; it turns out that $FC(G)$ is a characteristic subgroup of G . Obviously, G is *icc* if and only if $FC(G) = 1$.

Theorem 1. *Let $G \neq 1$ be a group defined by an extension of an abelian group :*

$$1 \longrightarrow K \text{ abelian} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

*and $\theta : Q \longrightarrow \text{Aut}(K)$ the associated homomorphism. Then G is *icc* if and only if both :*

- (i) $K \setminus 1$ contains only infinite $\theta(Q)$ -orbits,
- (ii) the induced homomorphism $\theta : FC(Q) \longrightarrow \text{Aut}(K)$ is injective.

On the opposite, we consider the case of extensions of centerless groups. For any extension G of a group K we consider the homomorphism $\pi : G \longrightarrow \text{Aut}(K)$ defined by, for all $g \in G$, $k \in K$, $\pi(g)(k) = k^g$.

Theorem 2. *Let $G \neq 1$ be a group defined by an extension of a centerless group :*

$$1 \longrightarrow K \text{ centerless} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

*and $\theta : Q \longrightarrow \text{Out}(K)$ the associated homomorphism. Then G is *icc* if and only if both :*

- (i) K does not contain a non trivial normal subgroup N either finite or $\mathbb{Z}^n$ which is preserved by $\pi(G)$ and contains only finite $\pi(G)$ -orbits,
- (ii) the induced homomorphism $\theta : FC(Q) \longrightarrow \text{Out}(K)$ is injective.

¹Centre de Recherche de l'Armée de l'air, Ecole de l'air, F-13661 Salon de Provence air

²Centre de Mathématiques et d'Informatique, Université de Provence, 39 rue F.Joliot-Curie, F-13453 marseille cedex 13

E-mail : preaux@cmi.univ-mrs.fr

Mathematical subject classification : 20E45 , 20E22

Since in condition (i) N necessarily lies in $FC(K)$, and since an icc group is centerless, it follows immediatly :

Theorem 3. *Let G be a group defined by an extension of an icc group :*

$$1 \longrightarrow K_{icc} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

and $\theta : Q \longrightarrow Out(K)$ the associated homomorphism. Then G is icc if and only if the induced homomorphism $\theta : FC(Q) \longrightarrow Out(K)$ is injective.

Together with a few observations on word hyperbolic groups, one obtains as another corollary the following characterization of icc property for extensions of word hyperbolic groups.

Theorem 4. *Let G be defined by an extension of a non trivial word hyperbolic group :*

$$1 \longrightarrow K_{hyperbolic} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

and $\theta : Q \longrightarrow Out(K)$ the associated homomorphism. Then G is icc if and only if both :

- (i) K is icc,
- (ii) the induced homomorphism $\theta : FC(Q) \longrightarrow Out(K)$ is injective.

1. PROOF OF THE RESULTS.

Let G be a group, H a non empty subset of G , and $u, g \in G$. We use the notations $u^g = g^{-1}ug$ and $u^H = \{u^g | g \in H\}$, so that g^G denotes the conjugacy class of g in G ; $Z_G(g)$ stands for the centralizer of g in G and $Z(G)$ for the center of G . It follows immediatly that g^G is finite if and only if $Z_G(g)$ has a finite index in G . Let $\rho : G \longrightarrow Q$ denote the projection given by the extension.

Proof of theorem 1. Note that that the conjugacy class in G of $k \in K$, coincide with its $\theta(Q)$ -orbit. Obviously G icc implies that both conditions (i) and (ii) are satisfied. We prove the converse by showing that if G is not icc then either condition (i) or condition (ii) is not satisfied. So suppose that G is not icc ; since $G \neq 1$ there exists $u \neq 1$ in G such that its conjugacy class u^G in G is finite. If $u \in K$; then condition (i) is not satisfied, so suppose in the following that $u \in G \setminus K$. Then $\rho(u)$ lies in $FC(Q)$, for $\rho(u)^Q = \rho(u^G)$, and $\rho(u) \neq 1$. Let $k \in K$, then $[u, k] \in K$ has a finite conjugacy class in G , for $Z_G(u)$ and $Z_G(kuk^{-1})$ both have a finite index in G and their intersection lies in $Z_G([u, k])$. So that either, for all $k \in K$, $[u, k] = 1$, which implies that $\pi(u) = \theta(\rho(u)) = Id_K$ and then $\theta : FC(Q) \longrightarrow Aut(K)$ is non injective, or as above, condition (i) is not satisfied. $\square$

Proof of theorem 2. Suppose that G is not icc ; since $G \neq 1$ there exists $u \neq 1$ in G such that u^G is finite. If u lies in K , let N' be the subgroup of K generated by u^G ; N' is a non trivial subgroup of $FC(K)$ preserved by $\pi(G)$ and with only finite $\pi(G)$ -orbits. Moreover N' is a finitely generated FC -group and hence a finite central extension of a f.g. abelian group (cf. [Ne]) ; it follows that the set $Tor(N')$ of torsion elements of N' is a characteristic finite subgroup of N' and $N'/Tor(N')$ is $\mathbb{Z}^n$. Hence condition (i) is not satisfied by picking $N = Tor(N')$, if $Tor(N') \neq 1$, and $N = N'$ otherwise. If u lies in $G \setminus K$, then $\rho(u)$ is a non trivial element of $FC(Q)$. For any $k \in Q$, $[u, k]$ lies in K and has a finite conjugacy class in G , so that, as in the proof of theorem 1, we can suppose that $\pi(u)$ is the identity, and hence $\theta(\rho(u)) = 1$, for otherwise condition (i) is not satisfied. It follows that $\theta : FC(Q) \longrightarrow Out(K)$ is non injective, so that condition (ii) is not satisfied. We have proved that if both condition (i) and (ii) are satisfied then G is icc ; we now prove the converse. If K contains a non trivial subgroup preserved by $\pi(G)$ and with only finite $\pi(G)$ -orbits then each element of N has a finite conjugacy class in G and then G is not icc. If $\theta : FC(Q) \longrightarrow Out(K)$ is non injective, let $q \neq 1$ in $FC(Q)$ be such that $\theta(q) = 1$. Then there exists a lift $\bar{q}$ of q in G such that $\pi(\bar{q})$ is the identity of K , so that

$K \subset Z_G(\bar{q})$. For any $q_1 \in Z_Q(q)$, if $\bar{q}_1$ denotes a lift of q_1 in G , then $\bar{q}_1 \bar{q} \bar{q}_1^{-1} = \bar{q}k$ for some $k \in K$, which must belong to $Z(K)$ since $\pi(\bar{q}_1 \bar{q} \bar{q}_1^{-1}) = \pi(k)$ is the identity on K . Since $Z(K) = 1$, $Z_G(\bar{q})$ contains $\rho^{-1}(Z_Q(q))$, and hence has a finite index in G , so that G is not icc. $\square$

Proof of theorem 4. If K is a non trivial elementary group then K is not icc, and neither G , for $\text{Aut}(K)$ is finite. So that the assumption in theorem 4 is satisfied and we suppose in the following that K is non elementary. A non elementary word hyperbolic has a finite center (cf. [Gr]). It follows that each element of $Z(K)$ has a finite conjugacy class both in K and in G . Then, if $Z(K) \neq 1$ neither K nor G are icc so that the assumption of theorem 4 is satisfied. So suppose in the following that K is centerless, so that theorem 2 applies. In a non elementary word hyperbolic group K , $FC(K)$ turns out to be a finite characteristic subgroup of K . Hence theorem 2 shows that G is icc if and only both $\theta : FC(Q) \longrightarrow \text{Out}(K)$ is injective and $FC(K) = 1$. $\square$

REFERENCES

- [Co] Y.DE CORNULIER, *Infinite conjugacy classes in groups acting on trees*, preprint (2005).
- [Gr] M.GROMOV, *Hyperbolic groups* in "Essays in group theory", MSRI Publications, Springer (1987), 75–263.
- [HP] P.DE LA HARPE et J.-P.PRÉAUX, *Groupes fondamentaux des variétés de dimension 3 et algèbres d'opérateurs*, preprint arXiv:math.GR/0509449 v1 (2005).
- [ROIV] F.J.MURRAY et J.VON NEUMANN, *On rings of operators*, IV, Annals of Math. **44** (1943), 716–808.
- [Ne] W.NEUMANN, *Groups with only finite conjugacy classes*, Proc. of the L.M.S. **1** (1951), 178–187.
- [P1] J.-P.PRÉAUX, *Wreath product of groups with infinite conjugacy classes*, preprint (2006), arXiv:math.GR/0612685.
- [P2] J.-P.PRÉAUX, *Finite extension of group with infinite conjugacy classes*, preprint (2007), arXiv:math.GR/0703314.
- [P3] J.-P.PRÉAUX, *Split extensions of group with infinite conjugacy classes*, preprint (2007).